

ESC CONGRESS 2016

Doctor Al'bina Shapieva (EUD ID : 504556)

Russian National Center Of Surgery

Electrophysiology Department

2 Abrikosovsky Per.,

119991 - Moscow Russian Federation

Phone : +7 9067236371 -

Email : shapieva@gmail.com

Title : Long-Term Effects of Septal Myectomy on Sudden Cardiac Death in Patients with Hypertrophic Cardiomyopathy

Topic : 01.09 - Sudden death / resuscitation

Category : Bedside

Option : Young Investigator Award (YIA) Clinical Science

A. Shapieva¹, S. Dzemeshevich², D. Podolyak¹, A. Tsyganov¹ - (1) Russian Research Center of Surgery, Electrophysiology Department, Moscow, Russian Federation (2) Russian Research Center of Surgery, Myocardial dysfunction, Moscow, Russian Federation

Purpose. The aim of this study was to determine the long-term outcomes (all-cause mortality, sudden cardiac death (SCD), and incidence of appropriate and inappropriate implantable cardioverter defibrillator (ICD) therapy) after extensive left ventricular septal myectomy in patients with diffuse-generalized form of obstructive hypertrophic cardiomyopathy (HOCM).

Method. This study included 54 consecutive patients with diffuse-generalized form of HOCM treated with either the extensive left ventricular septal myectomy (group A) or medical therapy (group B). Group A consisted of 22 patients (50.1±11.8 y.o., 11 females, 1 LEOPARD syndrome) and group B consisted of 32 patients (43.9±15.1 y.o., 16 females, 1 Danon disease). All patients underwent ICD implantation. The risk of SCD was assessed for each patient in group A before and 1 year after surgery and in group B before ICD implantation using standard "HCM Risk-SCD" calculator.

Results. The mean follow-up period was 2.5±2.2 years. Risk of SCD before and after procedure amounted to 4.0±1.9 and 1.6±0.6 in group A and 4.4±2.8 in group B, respectively. At 1 year after surgery in group A thickness of interventricular septum and the left atrium decreased from 19±3 to 15±2 mm and 45±4 to 42±3 mm, respectively. The peak systolic pressure gradient in the outflow tract of the left ventricle decreased from 76±6 to 20±3 mm Hg. During the observation period 1 patient with LEOPARD syndrome died probably from electrical storm (group A) and 1 patient died from unknown cause (group B). The overall mortality was 4.5% in group A and 3.1% in group B (p=0.08). Appropriate ICD therapy was lower (4.5% vs 6.3%; p=0.04), but inappropriate ICD therapy was higher (9.1% vs 3.2%; p=0.03) in group A. All episodes of inappropriate ICD therapy were caused by very fast conducted atrial fibrillation.

Conclusions. Patients with diffuse-generalized form of HOCM who are treated with extensive left ventricular septal myectomy have good survival and low SCD risk, similar to that of patients with non-obstructive HCM. In addition patients in surgery group had an increased incidence of inappropriate ICD therapy.